

## Accepted Manuscript

Title: Cannabidiol, neuroprotection and neuropsychiatric disorders

Author: Alline C. Campos Manoela V. Fogaça Andreza B. Sonogo Francisco S. Guimarães



PII: S1043-6618(16)00039-6  
DOI: <http://dx.doi.org/doi:10.1016/j.phrs.2016.01.033>  
Reference: YPHRS 3059

To appear in: *Pharmacological Research*

Received date: 1-11-2015  
Revised date: 29-1-2016  
Accepted date: 29-1-2016

Please cite this article as: Campos Alline C, Fogaça Manoela V, Sonogo Andreza B, Guimarães Francisco S. Cannabidiol, neuroprotection and neuropsychiatric disorders. *Pharmacological Research* <http://dx.doi.org/10.1016/j.phrs.2016.01.033>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

## Title page

## Cannabidiol, neuroprotection and neuropsychiatric disorders

Alline C Campos<sup>1,2,\*</sup>; Manoela V Fogaça<sup>1,2</sup>; Andreza B. Sonego<sup>1,2</sup>; Francisco S Guimarães<sup>1,2</sup>


<sup>1</sup> Department of Pharmacology, Medical School of of Ribeirão Preto, University of São Paulo, Bandeirantes Avenue, 3900, 14049-900, Ribeirão Preto, São Paulo, Brazil.

<sup>2</sup> Center of Interdisciplinary Research on Applied Neurosciences (NAPNA), University of São Paulo, Brazil.

\*Corresponding author

Alline C. de Campos, B.Pharm, M.Sc., Ph.D.  
Department of Pharmacology  
School of Medicine of Ribeirão Preto  
University of São Paulo  
3900 Bandeirantes avenue  
Ribeirão Preto-SP, Brazil  
Phone: +551633153325

Graphical abstract



**Abstract**

Cannabidiol (CBD) is a non-psychotomimetic phytocannabinoid derived from *Cannabis sativa*. It has possible therapeutic effects over a broad range of neuropsychiatric disorders. CBD attenuates brain damage associated with neurodegenerative and/or ischemic conditions. It also has positive effects on attenuating psychotic-, anxiety- and depressive-like behaviors. Moreover, CBD affects synaptic plasticity and facilitates neurogenesis. The mechanisms of these effects are still not entirely clear but seem to involve multiple pharmacological targets. In the present review, we summarized the main biochemical and molecular mechanisms that have been associated with the therapeutic effects of CBD, focusing on their relevance to brain function, neuroprotection and neuropsychiatric disorders.

**Keywords:** neuropsychiatric disorders; cannabidiol; oxidative stress; endocannabinoids; 5HT1A receptor; PPAR- $\gamma$  receptor

## 1. Introduction

Neuropsychiatric disorders are complex medical conditions that affect millions of people worldwide, being one of the most common causes of incapacity [1]. Although these disorders are caused by a complex interaction of several factors, such as genes and the environment [3,4], their specific etiology remains poorly understood. Consequently, patients still have a limited access to effective treatments [4].

In last decade, the research in this area has focused on the neuroplastic cellular processes responsible for brain adaptation as new therapeutic targets for these disorders [5-7]. Common features involving neuroprotective mechanisms (oxidative stress, immune mediators, neurotrophic factors) have been described, together with high levels of comorbidity between apparently different disorders [8-10]. As a possible reflect of these common features, the therapeutic indications of traditional treatments have also expanded. Antidepressants, for example, are the first-line treatment for depression and some anxiety disorders. However, they also possess neuroprotective properties, preventing the formation of amyloid plaques in a transgenic mouse model and humans and having positive effects in stroke [12]. The mechanisms underlying these effects involve multiple targets, such as elevation of brain-derived neurotrophic factor (BDNF) levels [13, 14], reduction of microglia activation, and decreased levels of proinflammatory mediators [15, 16].

Cannabinoids have also emerged as a new class of drugs with potential effects over a broad range of neurodegenerative and psychiatric disorders [17, 18]. The term cannabinoids refer to a heterogeneous group of compounds classified into three main groups: endogenous, synthetic and phytocannabinoids [17, 19]. Phytocannabinoids consist of terpenophenolic substances derived from the *Cannabis sativa* plant. The plant produces at least 66 compounds, including  $\Delta^9$ -tetrahydrocannabinol (THC), the one

responsible for its main psychological effects, whereas cannabidiol (CBD) is the major non-psychotomimetic compound present in the plant [20, 21].

The investigation of the possible positive impact of CBD in neuropsychiatric disorders began in the 1970s. After a slow progress, this subject has been showing an exponential growth in the last decade (Figure 1). CBD exhibits a broad spectrum of potential therapeutic properties in animal models and humans, including anxiolytic [17, 22], antidepressant [23], neuroprotective [17, 24-28], anti-inflammatory [29-32], and immunomodulatory [33, 34]. Regarding the latter, CBD decreases the production of inflammatory cytokines, the activation of microglial cells [31, 35, 36], and brain leucocytes infiltration in experimental autoimmune encephalitis [35]. Moreover, treatment based on this phytocannabinoid preserves cerebral circulation during ischemic events and reduces vascular changes and neuroinflammation in a model of sepsis-related encephalitis [26, 36-38].

Several clinical trials using CBD alone or in combination with other cannabinoids are under development. For example, the drug is being tested for Schizophrenia and cognitive dysfunction related to Schizophrenia (phase 2), Huntington's Disease (phase 2), and multiple sclerosis (MS-phase 3). Of note, the GW compound Sativex® showed to provide positive effects for the relief of MS-related spasticity while Epidolex® is currently in phase 3 trial for the treatment of orphan pediatric epilepsy syndrome [39]. CBD also has a better safety profile compared to other cannabinoids, such as THC. For instance, high doses of CBD (up to 1,500 mg/day) are well tolerated in animals and humans [40]. In addition, it does not change heart rate, blood pressure or body temperature, does not induce catalepsy, and does not alter psychomotor or psychological functions like THC [40]. This improved safety profile is probably reflecting its lack of direct agonist properties at cannabinoid receptors [41].

The mechanisms responsible for the wide range of CBD potential neuroprotective effects in neuropsychiatric disorders are not completely understood. New findings obtained in the last decade indicate that they involve multiple pharmacological targets. In the present review, we tried to summarize and discuss the importance of the main targets that have been associated with CBD neuroprotective action (table 1) and its effects on neuropsychiatric disorders.

### *1.1- The endocannabinoid system*

The endocannabinoid system comprises mainly the cannabinoid CB1 and CB2 receptors, their endogenous agonists, the endocannabinoids (ECs) anandamide (AEA) and 2-arachydonoilglycerol (2-AG), and the proteins responsible for their uptake, synthesis, and degradation. Cannabinoid receptors are coupled to a Gi/o protein and, once activated, increase K<sup>+</sup> cell influx leading to membrane hyperpolarization [42]. As a consequence, the probability of neurotransmitter release from the pre-synaptic terminal decreases, characterizing the ECs as retrograde messengers [43].

Several cannabinoid compounds have been shown to induce neuroprotection by acting on CB1 and/or CB2 receptors [44-46]. Although numerous *in vitro* studies suggest that CBD has a very low affinity for CB1 and CB2 receptors [47, 48], some of the effects of this drug seem to involve these receptors. This apparent contradiction could be explained by an *in vivo* drug action as an antagonist/inverse agonist at cannabinoid receptors or, more probably, by an indirect increase of anandamide levels through inhibition of its metabolism/uptake [49-52]. Corroborating the latter possibility, the CB1 receptor inverse agonist AM251 blocked CBD effects on both extinction and reconsolidation of conditioned fear [53], and on mice marble burying behavior [54].

In a model of newborn hypoxic-ischemic brain damage CBD, *in vivo* and *in vitro*, prevented the decrease in the number of viable neurons and attenuated the increase in

excitotoxicity, oxidative stress and inflammation (Table 1) [51, 55]. In brain slices submitted to hypoxic conditions, CBD effects on the production of IL-6, TNF- $\alpha$ , and COX-2 induction were attenuated by AM630, a CB2 antagonist [51]. This response, interestingly, did not depend on a CBD-induced increase in ECs levels [51]. In another study, CBD reduced  $\beta$ -amyloid-induced microglial activation *in vitro*, a model used to study some of the alterations found in Alzheimer's disease. It also decreased LPS-induced nitrite generation. Although the first effect depended on CB1 and CB2 receptor, the latter was not affected by pre-treatment with cannabinoid antagonists [56]. Besides CBD alone, its 1:1 combination with  $\Delta(9)$ -tetrahydrocannabinol (THC) in the phytocannabinoid-based medicine Sativex® also produced neuroprotective effects through a CB1- and CB2-mediated mechanism in a model of Huntington's disease [57]. CBD might also facilitate the survival of newborn hippocampal neurons via facilitation of anandamide neurotransmission through CB1 receptors [52, 58].

### 1.2- 5HT1A receptors

Seven different types of serotonin receptors have been identified so far: one ionotropic and six G-protein coupled. The 5HT1 class is coupled to a Gi/o protein and includes five subtypes: 5HT1A, 5HT1B, 5HT1D, 5HT1E, and 5HT1F. From this family, the 5HT1A is the main receptor related to CBD neuroprotective effects [18]. These receptors are present in pre-synaptic membranes as autosomic receptors and also found post-synaptically in several brain areas [59].

A pioneer *in vitro* study by Russo and colleagues [60] suggested that CBD could facilitate 5HT1A-mediated neurotransmission by acting as an agonist at these receptors [41]. Following this initial study, our group confirmed that the acute anxiolytic effects of CBD depend on facilitation of 5HT1A-mediated neurotransmission [17, 61]. Recent



findings, however, indicate that CBD is not a 5HT1A receptor agonist as originally proposed. Although not yet clear, its 5HT1A-mediated effects could involve allosteric interactions with the receptor binding site and/or interference with intracellular pathways [62, 63].

Regarding CBD effects on models of neuropsychiatry disorders, CBD peripheral injections attenuated acute autonomic responses evoked by stress, induced anxiolytic and panicolytic-like effects after intra-dorsal periaqueductal gray injections by activating 5HT1A receptors [61, 64-66]. The same 5HT1A receptor-dependent mechanism was observed after CBD injections into the bed nucleus of stria terminalis and prefrontal cortex [67-69]. CBD acute or chronic peripheral injections also induced antidepressant-like effects by activation of 5HT1A receptors [23, 20].

Part of the neuroprotective effects induced by CBD has also been associated with 5HT1A-mediated mechanisms. Pretreatment with the 5HT1A receptor antagonist WAY100635 prevented CBD reduction of brain tissue damage caused by cerebral artery occlusion [38, 71]. Furthermore, Magen and colleagues [72] suggested that 5HT1A receptors mediate CBD positive effects on the cognitive and locomotor deficits observed in a model of encephalopathy in mice, an effect replicated in another model using thioacetamide-induced liver failure [73]. More recently, Pazos and colleagues [55] showed that prevention of hypoxic-induced brain damage by CBD is not only mediated by 5HT1A, but also by CB2 receptors. Moreover, it is possible that the formation of heterodimers between these two receptors could account for CBD effects [55].

### *1.3- Oxidative stress and Peroxisome Proliferator-Activated Receptor gamma (PPAR $\gamma$ )*

Oxidative stress is characterized by an imbalance between the production of reactive oxygen/nitrogen species (ROS/RNS) and antioxidative protection systems in favor of

the oxidant species. An exacerbated production of ROS/RNS can be harmful to the body, since these compounds are highly reactive, with the biomolecules leading to the peroxidation of the polyunsaturated fatty acids, nitration and carbonylation of proteins and oxidation of DNA, leading ultimately to cellular death [74, 75]. In addition to excessive ROS/RNS generation, a reduced activity of the antioxidant system, including its enzymatic components such as superoxide dismutase (SOD), catalase (CAT) and glutathione peroxidase (GPx), and non-enzymatic ones, such as glutathione (GSH), are also observed [74, 75].

CBD has antioxidant properties that depend on its chemical structure, showing neuroprotective effects by decreasing oxidative parameters and increasing cell viability. It can donate electrons under a variable voltage potential as well as prevent dihydrorhodamine oxidation in the Fenton reaction similarly to the antioxidant butylhydroxytoluene (BHT). Also, CBD protects, in a concentration-related manner, neurons incubated with tert-butyl hydroperoxide. Moreover, it is significantly more effective than other antioxidants ( $\alpha$ -tocopherol and ascorbate) in a glutamate toxicity model [76]. Confirming these results, CBD attenuated tyrosine nitration, an indirect measure of the formation of ONOO<sup>-</sup>, and reduced apoptosis of retinal neurons in rats subjected to intravitreal injection of NMDA [77]. In a diabetic retinopathy model, CBD reduced tyrosine nitration and malondialdehyde (MDA) levels, a measure of lipid peroxidation, as well as decreased neural cell death [78]. These effects were also observed in in vitro models of Alzheimer's disease and multiple sclerosis. CBD pretreatment reduced ROS accumulation, lipid peroxidation, caspase-3 levels and DNA fragmentation in PC12 cells stimulated by  $\beta$ -amyloid [79]. Regarding the culture of oligodendrocyte progenitor cells, CBD reduced ROS production and cell death induced by H<sub>2</sub>O<sub>2</sub> [80]. In contrast to these results, Massi et al. [81] observed that CBD had anti-

proliferative effects in human glioma cells by producing ROS and depleting GSH stocks. However, in non-transformed glial cells, the drug did not induce oxidative stress. Other cannabinoids such as anandamide behave in the same way [82]. Thus, the antioxidant or pro-oxidant properties of cannabinoids seem to depend on biochemical and cellular features of tumor versus non-tumor cells, such as differences in signal transduction and/or redox state [81].

Besides the direct effects of CBD on the production of ROS/RNS, this phytocannabinoid also increases the expression of components of antioxidative systems. For example, CBD up-regulated SOD mRNA levels in the substantia nigra of rats unilaterally lesioned with 6-hydroxydopamine, a Parkinson's disease model [44], and in the caudate-putamen of rats treated with 3-nitropropionic acid, a Huntington's disease model [83]. In addition to ameliorating oxidative stress by acting as a scavenger of oxidant species [76], CBD could also act, at least in part, through receptor-dependent mechanisms, such as the peroxisome proliferator-activated receptor gamma (PPAR $\gamma$ ) [18, 84]. PPARs are a nuclear hormone receptors family that have their activities regulated by steroids and metabolites derived from lipid. So far, three different PPAR isoforms (PPAR $\alpha$ , PPAR $\beta$ , also called  $\delta$ , and PPAR $\gamma$ ) have been described [85]. Several pieces of evidence suggested that PPAR $\gamma$  receptors could be an attractive drug target for inflammatory-associated neuropsychiatric disorders, including neurodegenerative diseases [29, 86, 87]. PPAR $\gamma$  receptors seem to be involved in cellular proliferation, apoptosis and reduction of damage induced by ROS. Its activation inhibits the transcription of pro-inflammatory genes, preventing the NF- $\kappa$ B signaling pathway [86, 87].

CBD prevents amyloid- $\beta$ -induced neuronal death by its ability to scavenge ROS [88] and reduce oxidative stress. PPAR $\gamma$  seems to be relevant for these effects by interacting

with the transcription factor nuclear factor-erythroid 2-related Factor 2 (Nrf-2) [18]. Nrf-2 and PPAR $\gamma$  regulate each other. There are binding sites for Nrf-2 (AREs, antioxidant response elements) in the PPAR $\gamma$  promoter and PPAR responsive elements (PPERs) in the Nrf-2 promoter [89]. Moreover, gene expression associated with oxidative stress is controlled by Nrf-2 [90]. Recently, we have found that CBD prevented microglial activation by LPS *in vitro* by activating PPAR $\gamma$ , an effect that was associated with impairment of the NF- $\kappa$ B pathway [91].

In addition, using a murine genetic model of Alzheimer's Disease, Esposito and coworkers [29] suggested the PPAR $\gamma$ -mediated effect of CBD could also involve the decrease of activated glial cells and neuronal death, and facilitation of hippocampal neurogenesis. Recently, the same group showed that CBD increased neuronal survival by reducing apoptosis and decreasing amyloid precursor protein levels through activation of PPAR $\gamma$  receptors [92].

#### *1.4- Immune mediators, BDNF, and other related mechanisms*

The beneficial effects of CBD on brain disorders have also been associated with its capacity of modulating pro-inflammatory cytokines and BDNF expression, and interfering with intracellular pathways involved in neuronal fate [17, 18]. In a model of hepatic encephalopathy, CBD chronic treatment ameliorates cognitive and locomotor activity by restoring brain BDNF levels and decreasing mRNA expression of the type-1 TNF- $\alpha$  receptors [37]. In another study, using an intravital microscopy technique, CBD decreased leucocyte migration to the central nervous system and TNF- $\alpha$  expression induced by the previous administration of LPS [37]. Lower brain levels of BDNF and increased pro-inflammatory cytokines were correlated with poor cognitive performance in rats submitted to an experimental model of meningitis. These effects were attenuated by CBD treatment [93]. CBD also increased BDNF levels in the

hippocampus of rats subjected to a model of amphetamine-induced oxidative stress, a proposed model to study mania [94]. Recently, Campos and colleagues [24] suggested that the neuroprotective effects of CBD in a murine model of cerebral malaria are associated with its anti-inflammatory activity (by decreasing TNF $\alpha$  and IL-6 levels in the prefrontal cortex and hippocampus) and its capacity for up-regulating BDNF expression in the hippocampus.

In addition, CBD decreased microglia activation in murine models of Alzheimer's disease and schizophrenia [30, 95]. In experimental autoimmune encephalomyelitis (EAE), a murine model that mimics some aspects of multiple sclerosis, CBD decreased microglial cells activation by regulating STAT1/STAT3 balance and Th17 proliferation and function [35, 90]. CBD also decreased IL-6 and IL-17 release and reduced the severity of EAE [20]. Recently, Kozela and coworkers [96] suggested that CBD immunoregulatory effects rely on a strong up-regulation of inhibitory molecules on CD4<sup>+</sup>CD25<sup>-</sup> T cells.

During inflammatory conditions, the activation of mitogen-activated protein kinases, such as p38/MAP-kinase, might lead to the production of pro-inflammatory mediators. Thereby CBD, through its antioxidant properties, could inhibit the phosphorylation of p38, reducing the neurotoxic effects of an uncontrolled immune response [88]. CBD also reverts tau hyperphosphorylation through the blocked of the GSK3-beta pathway [97]. Moreover, in neural precursor cells cultures, this phytocannabinoid increased the expression, in a time-dependent manner, of the phosphorylated forms ERK1/2 and AKT. Given the positive effects of CBD on adult hippocampal neurogenesis [52, 15, 58], these observations suggest that the pro-proliferative and pro-survival effects of CBD involve a dynamic and complex process of recruiting these intracellular pathways. Accordingly, the neuroprotective effect of repeated CBD administration on brain

changes caused by chronic unpredictable stress (reduced neurogenesis and dendrite remodeling) or pilocarpine-induced seizures seem to be associated with a facilitation of autophagy, a mechanism essential for cell health [98, Hosseinzadeh et al., 2016]. This possible CBD mechanism could also be important for its antitumor effects [100].

### *1.5- Inhibition of adenosine uptake*

Enhancement of adenosine signaling, by inhibition of its uptake, has been proposed to mediate part of the anti-inflammatory, immunosuppressive, neuroprotective and behavioral effects of CBD [101]. Consistent with this proposal, CBD increased extracellular levels of adenosine [102]. Moreover, A2A receptor antagonists prevented CBD effects in a model of multiple sclerosis [30], the neuroprotection observed in brain slices of newborn mice after hypoxia and glucose deprivation [51], and rat microglial activation by LPS [103]. It failed, however, to block CBD neuroprotective effect against 3-nitropropionic acid [83]. Behaviorally, adenosine A1 or A2A receptor antagonists prevented CBD memory effects in adult zebrafish [104]. Indirect activation of A1 receptors also seems to be involved in CBD modulation of the pain pathways [105].

## **2.0- Conclusions and perspectives**

As for other effects of this remarkable phytocannabinoid [17, 113], the neuroprotective properties of CBD seem to depend on several cannabinoid-dependent and independent mechanisms (Figure 2). Despite the mechanisms involved, however, the preclinical evidence reviewed here, associated with the already reported safety profile of CBD in humans [22, 40], clearly indicate that CBD represents a new opportunity for the treatment of several brain disorders (such as neurodegenerative and neuropsychiatric) where neuronal loss or damage plays a significant role.

### 3.0- References

- [1] World Health Organization. WHO methods and data sources for global burden of disease estimates 2000-2011. Department of Health Statistics and Information Systems WHO, Geneva (2013).
- [2] M. Roy, M.G. Tapadia, S. Joshi, and B. Koch. Molecular and genetic basis of depression. *Journal of Genetics*. 93:879-892 (2014).
- [3] N. Lopizzo, L. Bocchio Chiavetto, N. Cattane, G. Plazzotta, F.I. Tarazi, C.M. Pariante, M.A. Riva, and A. Cattaneo. Gene-environment interaction in major depression: focus on experience-dependent biological systems. *Frontiers in Psychiatry*. 6:68 (2015).
- [4] M.R. Levinstein and B.A Samuels. Mechanisms underlying the antidepressant response and treatment resistance. *Frontiers in Behavioral Neuroscience*. 8: 208 (2014).
- [5] P.W. Kalivas and C. O'Brien. Drug addiction as a pathology of staged neuroplasticity. *Neuropsychopharmacology*. 33:166-180 (2008).
- [6] E. Castren. Neuronal network plasticity and recovery from depression. *JAMA Psychiatry*. 70:983-989 (2013).
- [7] D.S. Bredt, M.L. Furey, G. Chen, T. Lovenberg, W.C. Drevets, and H.K. Manji. Translating depression biomarkers for improved targeted therapies. *Neuroscience and Biobehavioral Reviews*. 59:1-15 (2015).
- [8] A. Shah, F.R. Carreno, and A. Frazer. Therapeutic modalities for treatment resistant depression: focus on vagal nerve stimulation and ketamine. *Clinical Psychopharmacology and Neuroscience*. 12:83-93 (2014).
- [9] K. Zhang, H. Jiang, Q. Zhang, J. Du, Y. Wang, and M. Zhao. Brain-derived neurotrophic factor serum levels in heroin-dependent patients after 26 weeks of withdrawal. *Comprehensive Psychiatry*. 65:150-155 (2016).
- [10] I.N. Wee, N. Kandiah, S. Acharyya, R.J. Chander, A. Ng, W.L. Au, and L.C. Tan. Depression and anxiety are co-morbid but dissociable in mild Parkinson's disease: A

prospective longitudinal study of patterns and predictors. *Parkinsonism & Related Disorders* (2015).

[11] Y.I. Sheline, T. West, K. Yarasheski, R. Swarm, M.S. Jasielec, J.R. Fisher, W.D. Ficker, P. Yan, C. Xiong, C. Frederiksen, M.V. Grzelak, R. Chott, R.J. Bateman, J.C. Morris, M.A. Mintun, J.M. Lee, and J.R. Cirrito. An antidepressant decreases CSF Abeta production in healthy individuals and in transgenic AD mice. *Science Translational Medicine*. 6:236re234 (2014).

[12] K.L. Ng, E.M. Gibson, R. Hubbard, J. Yang, B. Caffo, R.J. O'Brien, J.W. Krakauer, and S.R. Zeiler. Fluoxetine Maintains a State of Heightened Responsiveness to Motor Training Early After Stroke in a Mouse Model. *Stroke: a Journal of Cerebral Circulation*. 46:2951-2960 (2015).

[13] E.J. Nestler, M. Barrot, R.J. DiLeone, A.J. Eisch, S.J. Gold, and L.M. Monteggia. Neurobiology of depression. *Neuron*. 34:13-25 (2002).

[14] X. Xue, S. Shao, W. Wang, and F. Shao. Maternal separation induces alterations in reversal learning and brain-derived neurotrophic factor expression in adult rats. *Neuropsychobiology*. 68:243-249 (2013).

[15] D. Liu, Z. Wang, S. Liu, F. Wang, S. Zhao, and A. Hao. Anti-inflammatory effects of fluoxetine in lipopolysaccharide(LPS)-stimulated microglial cells. *Neuropharmacology*. 61:592-599 (2011).

[16] K. Ramirez, D.T. Shea, D.B. McKim, B.F. Reader, and J.F. Sheridan. Imipramine attenuates neuroinflammatory signaling and reverses stress-induced social avoidance. *Brain, Behavior, and Immunity*. 46:212-220 (2015).

[17] A.C. Campos, F.A. Moreira, F.V. Gomes, E.A. Del Bel, and F.S. Guimaraes. Multiple mechanisms involved in the large-spectrum therapeutic potential of cannabidiol in psychiatric disorders. *Philosophical Transactions of the Royal Society of London Series B, Biological Sciences*. 367:3364-3378 (2012).

[18] J. Fernandez-Ruiz, O. Sagredo, M.R. Pazos, C. Garcia, R. Pertwee, R. Mechoulam, and J. Martinez-Orgado. Cannabidiol for neurodegenerative disorders:



important new clinical applications for this phytocannabinoid? *British Journal of Clinical Pharmacology*. 75:323-333 (2013).

[19] E. Russo and G.W. Guy. A tale of two cannabinoids: the therapeutic rationale for combining tetrahydrocannabinol and cannabidiol. *Medical Hypotheses*. 66:234-246 (2006).

[20] R. Mechoulam and Y. Gaoni. Hashish. IV. The isolation and structure of cannabinolic cannabidiolic and cannabigerolic acids. *Tetrahedron*. 21:1223-1229 (1965).

[21] R.G. Pertwee. Pharmacological actions of cannabinoids. *Handbook of Experimental Pharmacology* :1-51 (2005).

[22] A.W. Zuardi, R.A. Cosme, F.G. Graeff, and F.S. Guimaraes. Effects of ipsapirone and cannabidiol on human experimental anxiety. *Journal of Psychopharmacology*. 7:82-88 (1993).

[23] R. Linge, L. Jimenez-Sanchez, L. Campa, F. Pilar-Cuellar, R. Vidal, A. Pazos, A. Adell, and A. Diaz. Cannabidiol induces rapid-acting antidepressant-like effects and enhances cortical 5-HT/glutamate neurotransmission: role of 5-HT receptors. *Neuropharmacology*. 103:16-26 (2015).

[24] A.C. Campos, F. Brant, A.S. Miranda, F.S. Machado, and A.L. Teixeira. Cannabidiol increases survival and promotes rescue of cognitive function in a murine model of cerebral malaria. *Neuroscience*. 289:166-180 (2015).

[25] J.W. Silveira, A.C. Issy, V.A. Castania, C.E. Salmon, M.H. Nogueira-Barbosa, F.S. Guimaraes, H.L. Defino, and E. Del Bel. Protective effects of cannabidiol on lesion-induced intervertebral disc degeneration. *PloS one*. 9:e113161 (2014).

[26] A.P. Schiavon, L.M. Soares, J.M. Bonato, H. Milani, F.S. Guimaraes, and R.M. Weffort de Oliveira. Protective effects of cannabidiol against hippocampal cell death and cognitive impairment induced by bilateral common carotid artery occlusion in mice. *Neurotoxicity Research*. 26:307-316 (2014).

- [27] M. Perez, S.U. Benitez, L.P. Cartarozzi, E. Del Bel, F.S. Guimaraes, and A.L. Oliveira. Neuroprotection and reduction of glial reaction by cannabidiol treatment after sciatic nerve transection in neonatal rats. *The European Journal of Neuroscience*. 38:3424-3434 (2013).
- [28] M. Kwiatkoski, F.S. Guimaraes, and E. Del-Bel. Cannabidiol-treated rats exhibited higher motor score after cryogenic spinal cord injury. *Neurotoxicity Research*. 21:271-280 (2012).
- [29] G. Esposito, C. Scuderi, M. Valenza, G.I. Togna, V. Latina, D. De Filippis, M. Cipriano, M.R. Carratu, T. Iuvone, and L. Steardo. Cannabidiol reduces Abeta-induced neuroinflammation and promotes hippocampal neurogenesis through PPARgamma involvement. *PloS one*. 6:e28668 (2011).
- [30] M. Mecha, A. Feliu, P.M. Inigo, L. Mestre, F.J. Carrillo-Salinas, and C. Guaza. Cannabidiol provides long-lasting protection against the deleterious effects of inflammation in a viral model of multiple sclerosis: a role for A2A receptors. *Neurobiology of Disease*. 59:141-150 (2013).
- [31] M.H. Napimoga, B.B. Benatti, F.O. Lima, P.M. Alves, A.C. Campos, D.R. Pena-Dos-Santos, F.P. Severino, F.Q. Cunha, and F.S. Guimaraes. Cannabidiol decreases bone resorption by inhibiting RANK/RANKL expression and pro-inflammatory cytokines during experimental periodontitis in rats. *International Immunopharmacology*. 9:216-222 (2009).
- [32] S. Ben-Shabat, L.O. Hanus, G. Katzavian, and R. Gallily. New cannabidiol derivatives: synthesis, binding to cannabinoid receptor, and evaluation of their antiinflammatory activity. *Journal of Medicinal Chemistry*. 49:1113-1117 (2006).
- [33] A.M. Malfait, R. Gallily, P.F. Sumariwalla, A.S. Malik, E. Andreacos, R. Mechoulam, and M. Feldmann. The nonpsychoactive cannabis constituent cannabidiol is an oral anti-arthritic therapeutic in murine collagen-induced arthritis. *Proceedings of the National Academy of Sciences of the United States of America*. 97:9561-9566 (2000).

- [34] E. Kozela, M. Pietr, A. Juknat, N. Rimmerman, R. Levy, and Z. Vogel. Cannabinoids Delta(9)-tetrahydrocannabinol and cannabidiol differentially inhibit the lipopolysaccharide-activated NF-kappaB and interferon-beta/STAT proinflammatory pathways in BV-2 microglial cells. *The Journal of Biological Chemistry*. 285:1616-1626 (2010).
- [35] E. Kozela, N. Lev, N. Kaushansky, R. Eilam, N. Rimmerman, R. Levy, A. Ben-Nun, A. Juknat, and Z. Vogel. Cannabidiol inhibits pathogenic T cells, decreases spinal microglial activation and ameliorates multiple sclerosis-like disease in C57BL/6 mice. *British Journal of Pharmacology*. 163:1507-1519 (2011).
- [36] F.J. Alvarez, H. Lafuente, M.C. Rey-Santano, V.E. Mielgo, E. Gastiasoro, M. Rueda, R.G. Pertwee, A.I. Castillo, J. Romero, and J. Martinez-Orgado. Neuroprotective effects of the nonpsychoactive cannabinoid cannabidiol in hypoxic-ischemic newborn piglets. *Pediatric Research*. 64:653-658 (2008).
- [37] L. Ruiz-Valdepenas, J.A. Martinez-Orgado, C. Benito, A. Millan, R.M. Tolon, and J. Romero. Cannabidiol reduces lipopolysaccharide-induced vascular changes and inflammation in the mouse brain: an intravital microscopy study. *Journal of Neuroinflammation*. 8:5 (2011).
- [38] K. Mishima, K. Hayakawa, K. Abe, T. Ikeda, N. Egashira, K. Iwasaki, and M. Fujiwara. Cannabidiol prevents cerebral infarction via a serotonergic 5-hydroxytryptamine1A receptor-dependent mechanism. *Stroke*. 36:1077-1082 (2005).
- [39] <https://clinicaltrials.gov> (accessed 01/2016).
- [40] M.M. Bergamaschi, R.H. Queiroz, A.W. Zuardi, and J.A. Crippa. Safety and side effects of cannabidiol, a Cannabis sativa constituent. *Current Drug Safety*. 6:237-249 (2011).
- [41] C. Scuderi, D.D. Filippis, T. Iuvone, A. Blasio, A. Steardo, and G. Esposito. Cannabidiol in medicine: a review of its therapeutic potential in CNS disorders. *Phytotherapy Research*. 23:597-602 (2009).
- [42] B. Szabo and E. Schlicker. Effects of cannabinoids on neurotransmission. *Handbook of Experimental Pharmacology*. 327-365 (2005).

- [43] A.C. Kreitzer. Neurotransmission: emerging roles of endocannabinoids. *Current Biology*. 15:R549-551 (2005).
- [44] M. Garcia-Arencibia, S. Gonzalez, E. de Lago, J.A. Ramos, R. Mechoulam, and J. Fernandez-Ruiz. Evaluation of the neuroprotective effect of cannabinoids in a rat model of Parkinson's disease: importance of antioxidant and cannabinoid receptor-independent properties. *Brain Research*. 1134:162-170 (2007).
- [45] M.R. Melis, S. Succu, M.S. Mascia, F. Sanna, T. Melis, M.P. Castelli, and A. Argiolas. The cannabinoid receptor antagonist SR-141716A induces penile erection in male rats: involvement of paraventricular glutamic acid and nitric oxide. *Neuropharmacology*. 50:219-228 (2006).
- [46] S. Kreutz, M. Koch, C. Bottger, C. Ghadban, H.W. Korf, and F. Dehghani. 2-Arachidonoylglycerol elicits neuroprotective effects on excitotoxically lesioned dentate gyrus granule cells via abnormal-cannabidiol-sensitive receptors on microglial cells. *Glia*. 57:286-294 (2009).
- [47] A. Thomas, R.A. Ross, B. Saha, A. Mahadevan, R.K. Razdan, and R.G. Pertwee. 6"-Azidohept-2"-yne-cannabidiol: a potential neutral, competitive cannabinoid CB1 receptor antagonist. *European Journal of Pharmacology*. 487:213-221 (2004).
- [48] A. Thomas, G.L. Baillie, A.M. Phillips, R.K. Razdan, R.A. Ross, and R.G. Pertwee. Cannabidiol displays unexpectedly high potency as an antagonist of CB1 and CB2 receptor agonists in vitro. *British Journal of Pharmacology*. 150:613-623 (2007).
- [49] T. Bisogno, L. Hanus, L. De Petrocellis, S. Tchilibon, D.E. Ponde, I. Brandi, A.S. Moriello, J.B. Davis, R. Mechoulam, and V. Di Marzo. Molecular targets for cannabidiol and its synthetic analogues: effect on vanilloid VR1 receptors and on the cellular uptake and enzymatic hydrolysis of anandamide. *British Journal of Pharmacology*. 134:845-852 (2001).
- [50] R.G. Pertwee. The diverse CB1 and CB2 receptor pharmacology of three plant cannabinoids: delta9-tetrahydrocannabinol, cannabidiol and delta9-tetrahydrocannabivarin. *British Journal of Pharmacology*. 153:199-215 (2008).

- [51] A. Castillo, M.R. Tolon, J. Fernandez-Ruiz, J. Romero, and J. Martinez-Orgado. The neuroprotective effect of cannabidiol in an in vitro model of newborn hypoxic-ischemic brain damage in mice is mediated by CB(2) and adenosine receptors. *Neurobiology of Disease*. 37:434-440 (2010).
- [52] A.C. Campos, Z. Ortega, J. Palazuelos, M.V. Fogaca, D.C. Aguiar, J. Diaz-Alonso, S. Ortega-Gutierrez, H. Vazquez-Villa, F.A. Moreira, M. Guzman, I. Galve-Roperh, and F.S. Guimaraes. The anxiolytic effect of cannabidiol on chronically stressed mice depends on hippocampal neurogenesis: involvement of the endocannabinoid system. *The international Journal of Neuropsychopharmacology*. 16:1407-1419 (2013).
- [53] C.A. Stern, L. Gazarini, R.N. Takahashi, F.S. Guimaraes, and L.J. Bertoglio. On disruption of fear memory by reconsolidation blockade: evidence from cannabidiol treatment. *Neuropsychopharmacology*. 37:2132-2142 (2012).
- [54] P.C. Casarotto, F.V. Gomes, L.B. Resstel, and F.S. Guimaraes. Cannabidiol inhibitory effect on marble-burying behaviour: involvement of CB1 receptors. *Behavioural Pharmacology*. 21:353-358 (2010).
- [55] M.R. Pazos, N. Mohammed, H. Lafuente, M. Santos, E. Martinez-Pinilla, E. Moreno, E. Valdizan, J. Romero, A. Pazos, R. Franco, C.J. Hillard, F.J. Alvarez, and J. Martinez-Orgado. Mechanisms of cannabidiol neuroprotection in hypoxic-ischemic newborn pigs: role of 5HT(1A) and CB2 receptors. *Neuropharmacology*. 71:282-291 (2013).
- [56] A.M. Martin-Moreno, D. Reigada, B.G. Ramirez, R. Mechoulam, N. Innamorato, A. Cuadrado, and M.L. de Ceballos. Cannabidiol and other cannabinoids reduce microglial activation in vitro and in vivo: relevance to Alzheimer's disease. *Molecular Pharmacology*. 79:964-973 (2011).
- [57] S. Valdeolivas, V. Satta, R.G. Pertwee, J. Fernandez-Ruiz, and O. Sagredo. Sativex-like combination of phytocannabinoids is neuroprotective in malonate-lesioned rats, an inflammatory model of Huntington's disease: role of CB1 and CB2 receptors. *ACS Chemical Neuroscience*. 3:400-406 (2012).

- [58] S.A. Wolf, A. Bick-Sander, K. Fabel, P. Leal-Galicia, S. Tauber, G. Ramirez-Rodriguez, A. Muller, A. Melnik, T.P. Waltinger, O. Ullrich, and G. Kempermann. Cannabinoid receptor CB1 mediates baseline and activity-induced survival of new neurons in adult hippocampal neurogenesis. *Cell Communication and Signaling*. 8:12 (2010).
- [59] L. Bevilacqua, P. Ardenghi, N. Schroder, E. Bromberg, P.K. Schmitz, E. Schaeffer, J. Quevedo, M. Bianchin, R. Walz, J.H. Medina, and I. Izquierdo. Drugs acting upon the cyclic adenosine monophosphate/protein kinase A signalling pathway modulate memory consolidation when given late after training into rat hippocampus but not amygdala. *Behavioural Pharmacology*. 8:331-338 (1997).
- [60] E.B. Russo, A. Burnett, B. Hall, and K.K. Parker. Agonistic properties of cannabidiol at 5-HT<sub>1A</sub> receptors. *Neurochemical Research*. 30:1037-1043 (2005).
- [61] A.C. Campos and F.S. Guimaraes. Involvement of 5HT<sub>1A</sub> receptors in the anxiolytic-like effects of cannabidiol injected into the dorsolateral periaqueductal gray of rats. *Psychopharmacology*. 199:223-230 (2008).
- [62] E.M. Rock, D. Bolognini, C.L. Limebeer, M.G. Cascio, S. Anavi-Goffer, P.J. Fletcher, R. Mechoulam, R.G. Pertwee, and L.A. Parker. Cannabidiol, a non-psychoactive component of cannabis, attenuates vomiting and nausea-like behaviour via indirect agonism of 5-HT<sub>1A</sub> somatodendritic autoreceptors in the dorsal raphe nucleus. *British Journal of Pharmacology*. 165:2620-2634 (2012).
- [63] E.M. Rock, J.M. Goodwin, C.L. Limebeer, A. Breuer, R.G. Pertwee, R. Mechoulam, and L.A. Parker. Interaction between non-psychoactive cannabinoids in marijuana: effect of cannabigerol (CBG) on the anti-nausea or anti-emetic effects of cannabidiol (CBD) in rats and shrews. *Psychopharmacology*. 215:505-512 (2011).
- [64] L.B. Resstel, R.F. Tavares, S.F. Lisboa, S.R. Joca, F.M. Correa, and F.S. Guimaraes. 5-HT<sub>1A</sub> receptors are involved in the cannabidiol-induced attenuation of behavioural and cardiovascular responses to acute restraint stress in rats. *British Journal of Pharmacology*. 156:181-188 (2009).

- [65] P. Soares V de, A.C. Campos, V.C. Bortoli, H. Zangrossi, Jr., F.S. Guimaraes, and A.W. Zuardi. Intra-dorsal periaqueductal gray administration of cannabidiol blocks panic-like response by activating 5-HT1A receptors. *Behavioural Brain Research*. 213:225-229 (2010).
- [66] A.C. Campos, V. de Paula Soares, M.C. Carvalho, F.R. Ferreira, M.A. Vicente, M.L. Brandao, A.W. Zuardi, H. Zangrossi, Jr., and F.S. Guimaraes. Involvement of serotonin-mediated neurotransmission in the dorsal periaqueductal gray matter on cannabidiol chronic effects in panic-like responses in rats. *Psychopharmacology*. 226:13-24 (2013).
- [67] F.V. Gomes, L.B. Resstel, and F.S. Guimaraes. The anxiolytic-like effects of cannabidiol injected into the bed nucleus of the stria terminalis are mediated by 5-HT1A receptors. *Psychopharmacology*. 213:465-473 (2011).
- [68] F.V. Gomes, D.G. Reis, F.H. Alves, F.M. Correa, F.S. Guimaraes, and L.B. Resstel. Cannabidiol injected into the bed nucleus of the stria terminalis reduces the expression of contextual fear conditioning via 5-HT1A receptors. *Journal of Psychopharmacology*. 26:104-113 (2012).
- [69] M.V. Fogaca, F.M. Reis, A.C. Campos, and F.S. Guimaraes. Effects of intra-prelimbic prefrontal cortex injection of cannabidiol on anxiety-like behavior: involvement of 5HT1A receptors and previous stressful experience. *European Neuropsychopharmacology*. 24:410-419 (2014).
- [70] T.V. Zanelati, C. Biojone, F.A. Moreira, F.S. Guimaraes, and S.R. Joca. Antidepressant-like effects of cannabidiol in mice: possible involvement of 5-HT1A receptors. *British Journal of Pharmacology*. 159:122-128 (2010).
- [71] K. Hayakawa, K. Mishima, M. Nozako, A. Ogata, M. Hazeckawa, A.X. Liu, M. Fujioka, K. Abe, N. Hasebe, N. Egashira, K. Iwasaki, and M. Fujiwara. Repeated treatment with cannabidiol but not Delta9-tetrahydrocannabinol has a neuroprotective effect without the development of tolerance. *Neuropharmacology*. 52:1079-1087 (2007).

- [72] I. Magen, Y. Avraham, Z. Ackerman, L. Vorobiev, R. Mechoulam, and E.M. Berry. Cannabidiol ameliorates cognitive and motor impairments in bile-duct ligated mice via 5-HT<sub>1A</sub> receptor activation. *British Journal of Pharmacology*. 159:950-957 (2010).
- [73] Y. Avraham, N. Grigoriadis, T. Poutahidis, L. Vorobiev, I. Magen, Y. Ilan, R. Mechoulam, and E. Berry. Cannabidiol improves brain and liver function in a fulminant hepatic failure-induced model of hepatic encephalopathy in mice. *British Journal of Pharmacology*. 162:1650-1658 (2011).
- [74] A.M. Pisoschi and A. Pop. The role of antioxidants in the chemistry of oxidative stress: A review. *European Journal of Medicinal Chemistry*. 97:55-74 (2015).
- [75] E. Niedzielska, I. Smaga, M. Gawlik, A. Moniczewski, P. Stankowicz, J. Pera, and M. Filip. Oxidative Stress in Neurodegenerative Diseases. *Molecular Neurobiology* (2015).
- [76] A.J. Hampson, M. Grimaldi, J. Axelrod, and D. Wink. Cannabidiol and (-)-Delta<sup>9</sup>-tetrahydrocannabinol are neuroprotective antioxidants. *Proceedings of the National Academy of Sciences of the United States of America*. 95:8268-8273 (1998).
- [77] A.B. El-Remessy, I.E. Khalil, S. Matragoon, G. Abou-Mohamed, N.J. Tsai, P. Roon, R.B. Caldwell, R.W. Caldwell, K. Green, and G.I. Liou. Neuroprotective effect of (-)-Delta<sup>9</sup>-tetrahydrocannabinol and cannabidiol in N-methyl-D-aspartate-induced retinal neurotoxicity: involvement of peroxynitrite. *The American Journal of Pathology*. 163:1997-2008 (2003).
- [78] A.B. El-Remessy, M. Al-Shabraway, Y. Khalifa, N.T. Tsai, R.B. Caldwell, and G.I. Liou. Neuroprotective and blood-retinal barrier-preserving effects of cannabidiol in experimental diabetes. *The American journal of pathology*. 168:235-244 (2006). [79] T. Iuvone, G. Esposito, R., Santamaria, M. Di Rosa, and A.A. Izzo. Neuroprotective effect of cannabidiol, a non-psychoactive component from *Cannabis sativa*, on beta-amyloid-induced toxicity in PC12 cells. *Journal of Neurochemistry*. 89:134-141 (2004).



- [80] M. Mecha, A.S. Torrao, L. Mestre, F.J. Carrillo-Salinas, R. Mechoulam, and C. Guaza. Cannabidiol protects oligodendrocyte progenitor cells from inflammation-induced apoptosis by attenuating endoplasmic reticulum stress. *Cell Death & Disease*. 3:e331 (2012).
- [81] P. Massi, A. Vaccani, S. Bianchessi, B. Costa, P. Macchi, and D. Parolaro. The non-psychoactive cannabidiol triggers caspase activation and oxidative stress in human glioma cells. *Cellular and Molecular Life Sciences*. 63:2057-2066 (2006).
- [82] K.P. Sarker, S. Obara, M. Nakata, I. Kitajima, and I. Maruyama. Anandamide induces apoptosis of PC-12 cells: involvement of superoxide and caspase-3. *FEBS Letters*. 472:39-44 (2000).
- [83] O. Sagredo, J.A. Ramos, A. Decio, R. Mechoulam, and J. Fernandez-Ruiz. Cannabidiol reduced the striatal atrophy caused 3-nitropropionic acid in vivo by mechanisms independent of the activation of cannabinoid, vanilloid TRPV1 and adenosine A2A receptors. *The European Journal of Neuroscience*. 26:843-851 (2007).
- [84] S.E. O'Sullivan. Cannabinoids go nuclear: evidence for activation of peroxisome proliferator-activated receptors. *British Journal of Pharmacology*. 152:576-582 (2007).
- [85] S. Kersten. Peroxisome proliferator activated receptors and lipoprotein metabolism. *PPAR Research*. 2008:132960 (2008).
- [86] L.C. Rodrigues, P.H. Gobira, A.C. de Oliveira, R. Pelicao, A.L. Teixeira, F.A. Moreira, and A.C. Campos. Neuroinflammation as a possible link between cannabinoids and addiction. *Acta Neuropsychiatrica*. 26:334-346 (2014).
- [87] P.F. Stahel, W.R. Smith, J. Bruchis, and C.H. Rabb. Peroxisome proliferator-activated receptors: "key" regulators of neuroinflammation after traumatic brain injury. *PPAR Research*. 2008:538141 (2008).
- [88] G. Esposito, D. De Filippis, M.C. Maiuri, D. De Stefano, R. Carnuccio, and T. Iuvone. Cannabidiol inhibits inducible nitric oxide synthase protein expression and nitric oxide production in beta-amyloid stimulated PC12 neurons through p38 MAP kinase and NF-kappaB involvement. *Neuroscience Letters*. 399:91-95 (2006).

- [89] I.H.Y. Cho, W. Gladwell, X. Wang, B. Chorley, D. Bell, S.P. Reddy, and S.R. Kleeberger. Nrf2-regulated PPAR $\gamma$  expression is critical to protection against acute lung injury in mice. *American journal of respiratory and critical care medicine*. 182:170-182 (2010).
- [90] A. Juknat, M. Pietr, E. Kozela, N. Rimmerman, R. Levy, G. Coppola, D. Geschwind, and Z. Vogel. Differential transcriptional profiles mediated by exposure to the cannabinoids cannabidiol and Delta9-tetrahydrocannabinol in BV-2 microglial cells. *British Journal of Pharmacology*. 165:2512-2528 (2012).
- [91] A. B. Sonogo, J. E. Sepulveda-Diaz, P. P. Michel, E. A. Del-Bel, F. S. Guimarães, R. Raisman-Vozar. Cannabidiol reduces LPS-induced activation and oxidative stress in primary microglial culture via PPAR $\gamma$  receptor. *Society for Neuroscience Annual Meeting*. Poster Abstract 4.10/R11 (2015).
- [92] C. Scuderi, L. Steardo, and G. Esposito. Cannabidiol promotes amyloid precursor protein ubiquitination and reduction of beta amyloid expression in SHSY5YAPP<sup>+</sup> cells through PPAR $\gamma$  involvement. *Phytotherapy Research*. 28:1007-1013 (2014).
- [93] T. Barichello, R.A. Ceretta, J.S. Generoso, A.P. Moreira, L.R. Simoes, C.M. Comim, J. Quevedo, M.C. Vilela, A.W. Zuardi, J.A. Crippa, and A.L. Teixeira. Cannabidiol reduces host immune response and prevents cognitive impairments in Wistar rats submitted to pneumococcal meningitis. *European Journal of Pharmacology*. 697:158-164 (2012).
- [94] S.S. Valvassori, G. Elias, B. de Souza, F. Petronilho, F. Dal-Pizzol, F. Kapczinski, C. Trzesniak, V. Tumas, S. Dursun, M.H. Chagas, J.E. Hallak, A.W. Zuardi, J. Quevedo, and J.A. Crippa. Effects of cannabidiol on amphetamine-induced oxidative stress generation in an animal model of mania. *Journal of Psychopharmacology*. 25:274-280 (2011).
- [95] F.V. Gomes, R. Llorente, E.A. Del Bel, M.P. Viveros, M. Lopez-Gallardo, and F.S. Guimaraes. Decreased glial reactivity could be involved in the antipsychotic-like effect of cannabidiol. *Schizophrenia Research*. 164:155-163 (2015).

- [96] E. Kozela, A. Juknat, N. Kaushansky, A. Ben-Nun, G. Coppola, and Z. Vogel. Cannabidiol, a non-psychoactive cannabinoid, leads to EGR2-dependent anergy in activated encephalitogenic T cells. *Journal of Neuroinflammation*. 12:52 (2015).
- [97] G. Esposito, C. Scuderi, C. Savani, L. Steardo, Jr., D. De Filippis, P. Cottone, T. Iuvone, V. Cuomo, and L. Steardo. Cannabidiol in vivo blunts beta-amyloid induced neuroinflammation by suppressing IL-1beta and iNOS expression. *British Journal of Pharmacology*. 151:1272-1279 (2007).
- [98] M.V. Fogaça, A.C. Campos, F.S. Guimarães. Behavioural effects of cannabidiol in chronically stressed mice is mediated by neurogenesis and autophagy through CB1 receptor activation. *European Neuropsychopharmacology*. 24 (Suppl 2):S606 (2014).
- [99] M. Hosseinzadeh, S. Nikseresht, F. Khodagholi, N. Naderi, and N. Maghsoudi. Cannabidiol Post-Treatment Alleviates Rat Epileptic-Related Behaviors and Activates Hippocampal Cell Autophagy Pathway Along with Antioxidant Defense in Chronic Phase of Pilocarpine-Induced Seizure. *Journal of Molecular Neuroscience*. (2016) *In press*.
- [100] M. Nabissi, M.B. Morelli, C. Amantini, S. Liberati, M. Santoni, L. Ricci-Vitiani, R. Pallini, and G. Santoni. Cannabidiol stimulates Aml-1a-dependent glial differentiation and inhibits glioma stem-like cells proliferation by inducing autophagy in a TRPV2-dependent manner. *International Journal of Cancer Journal international du Cancer*. 137:1855-1869 (2015).
- [101] E.J. Carrier, J.A. Auchampach, C.J. Hillard. Inhibition of an equilibrative nucleoside transporter by cannabidiol: a mechanism of cannabinoid immunosuppression. *Proceedings of the National Academy of Sciences U S A*. 103(20):7895-900 (2006).
- [102] S. Mijangos-Moreno, A. Poot-Ake, G. Arankowsky-Sandoval, and E. Murillo-Rodriguez. Intrahypothalamic injection of cannabidiol increases the extracellular levels of adenosine in nucleus accumbens in rats. *Neuroscience Research*. 84:60-63 (2014).
- [103] G.I. Liou, J.A. Auchampach, C.J. Hillard, G. Zhu, B. Yousufzai, S. Mian, S. Khan, Y. Khalifa. Mediation of cannabidiol anti-inflammation in the retina by

equilibrative nucleoside transporter and A2A adenosine receptor. *Investigative Ophthalmology & Visual Science*. 49(12):5526-31 (2008).

[104] L.R. Nazario, R. Antonioli Jr, K.M. Capiotti, J.E. Hallak, A.W. Zuardi, J.A. Crippa, C.D. Bonan, R.S. da Silva. Caffeine protects against memory loss induced by high and non-anxiolytic dose of cannabidiol in adult zebrafish (*Danio rerio*). *Pharmacology Biochemistry & Behavior*. 135:210-6 (2015).

[105] S. Maione, F. Piscitelli, L. Gatta, D. Vita, L. De Petrocellis, E. Palazzo, V. de Novellis, V. Di Marzo. Non-psychoactive cannabinoids modulate the descending pathway of antinociception in anaesthetized rats through several mechanisms of action. *British Journal of Pharmacology*. 162(3):584-96 (2011).

[106] K. Hayakawa, K. Mishima, K. Abe, N. Hasebe, F. Takamatsu, H. Yasuda, T. Ikeda, K. Inui, N. Egashira, K. Iwasaki, and M. Fujiwara. Cannabidiol prevents infarction via the non-CB1 cannabinoid receptor mechanism. *Neuroreport*. 15:2381-2385 (2004).

[107] I.K. Hayakawa, K. Mishima, M. Nozako, M. Hazekawa, K. Irie, M. Fujioka, K. Orito, K. Abe, N. Hasebe, N. Egashira, K. Iwasaki, and M. Fujiwara. Delayed treatment with cannabidiol has a cerebroprotective action via a cannabinoid receptor-independent myeloperoxidase-inhibiting mechanism. *Journal of Neurochemistry*. 102:1488-1496 (2007).

[108] I. Lastres-Becker, F. Molina-Holgado, J.A. Ramos, R. Mechoulam, and J. Fernandez-Ruiz. Cannabinoids provide neuroprotection against 6-hydroxydopamine toxicity in vivo and in vitro: relevance to Parkinson's disease. *Neurobiology of Disease*. 19:96-107 (2005).

[109] E. Janefjord, J.L. Maag, B.S. Harvey, and S.D. Smid. Cannabinoid effects on beta amyloid fibril and aggregate formation, neuronal and microglial-activated neurotoxicity in vitro. *Cellular and Molecular Neurobiology*. 34:31-42 (2014).


[110] B.S. Harvey, K.S. Ohlsson, J.L. Maag, I.F. Musgrave, and S.D. Smid. Contrasting protective effects of cannabinoids against oxidative stress and amyloid-beta evoked neurotoxicity in vitro. *Neurotoxicology*. 33:138-146 (2012).

- [111] G. Pryce, D.R. Riddall, D.L. Selwood, G. Giovannoni, and D. Baker. Neuroprotection in Experimental Autoimmune Encephalomyelitis and Progressive Multiple Sclerosis by Cannabis-Based Cannabinoids. *Journal of neuroimmune pharmacology*. 10:281-292 (2015).
- [112] S.J. Ward, S.D. McAllister, R. Kawamura, R. Murase, H. Neelakantan, and E.A. Walker. Cannabidiol inhibits paclitaxel-induced neuropathic pain through 5-HT(1A) receptors without diminishing nervous system function or chemotherapy efficacy. *British Journal of Pharmacology*. 171:636-645 (2014).
- [113] C. Ibeas Bih, T. Chen, A.V. Nunn, M. Bazelot, M. Dallas, and B.J. Whalley. Molecular Targets of Cannabidiol in Neurological Disorders. *Neurotherapeutics : the journal of the American Society for Experimental NeuroTherapeutics*. 12:699-730 (2015).
- [114] A.W. Zuardi, E. Finkelfarb, O.F. Bueno, R.E. Musty, and I.G. Karniol. Characteristics of the stimulus produced by the mixture of cannabidiol with delta 9-tetrahydrocannabinol. *Archives Internationales de Pharmacodynamie et de Therapie*. 249:137-146 (1981).
- [115] B. Hine, M. Torrelío, S. Gershon. Differential effect of cannabinol and cannabidiol on THC-induced responses during abstinence in morphine-dependent rats. *Res Communications Chemical Pathology Pharmacology*. 12(1):185-5 (1975).
- [116] I. Izquierdo, O.A. Orsingher, A.C. Berardi. Effect of cannabidiol and of other cannabis sativa compounds on hippocampal seizure discharges. *Psychopharmacologia*. 28(1):95-102 (1973).
- [117] P. Consroe, J. Laguna, J. Allender, S. Snider, L. Stern, R. Sandyk, K. Kennedy, K. Schram. Controlled clinical trial of cannabidiol in Huntington's disease. *Pharmacology Biochemistry & Behavior*. 40(3):701-8 (1991).
- [118] M. Moreno-Martet, F. Espejo-Porras, J. Fernández-Ruiz, E. de Lago. Changes in endocannabinoid receptors and enzymes in the spinal cord of SOD1(G93A) transgenic mice and evaluation of a Sativex(®) -like combination of phytocannabinoids: interest

for future therapies in amyotrophic lateral sclerosis. *CNS Neuroscience Therapeutics*. 20(9):809-15 (2014).


[119] D.T. Wade, P. Robson, H. House, P. Makela, J. Aram. A preliminary controlled study to determine whether whole-plant cannabis extracts can improve intractable neurogenic symptoms. *Clinical Rehabilitation*. 17(1):21-9 (2003).

**Figure 1.** Number of published papers in PubMed describing possible therapeutic effects of cannabidiol in neuropsychiatric disorders since 1970. The occasion of the first publication associating CBD effects with specific disorders is also displayed in the figure [references: 70, 114- 119]



**Figure 2.** Possible mechanisms responsible for the neuroprotective effects of cannabidiol





**Table 1** – Neuroprotective effects of CBD in different models and their suggested mechanisms of action

Model/Method	Treatment schedule	CBD dose/concentration range	CBD effect	Suggested mechanism of action
Newborn hypoxic-ischemic brain damage (HI)	15-min pre-incubation	0.1 to 1000 $\mu$ M	Reduced acute and apoptotic HI brain damage; Reduced glutamate, IL-6 concentration and TNF $\alpha$ , COX-2 and iNOS expression	CB <sub>2</sub> and A <sub>2A</sub> receptors
Newborn hypoxic-ischemic brain damage (HI)	30 min after HI	1 mg/kg, i.p.	Prevented the decrease in the number of viable neurons and the increase in excitotoxicity, oxidative stress and inflammation	CB <sub>2</sub> and 5HT <sub>1A</sub> receptors
Striatal lesions caused by 3-nitropropionic acid (3NP)	3NP (10 mg/kg, twice a day) and/or CBD injections for 5 days, i.p.	5 mg/kg, i.p.	Reversed 3NP-induced reductions in GABA contents and mRNA levels for substance P, neuronal-specific enolase and superoxide dismutase(SOD)-2; partially attenuated SOD-1 and proenkephalin mRNA.	Independent of CB <sub>1</sub> , TRPV <sub>1</sub> and A <sub>2A</sub> receptors
Middle cerebral artery (MCA) occlusion	Immediately before and 3 or 4h after MCA occlusion; 1 and 2h after reperfusion, i.p.	0.1, 1 and 3 mg/kg, i.p.	Suppressed the decrease in cerebral blood flow by the failure of cerebral microcirculation after reperfusion; inhibited myeloperoxidase (MPO) activity in neutrophils; reduced the number of MPO-immunopositive cells	Independent of CB <sub>1</sub> receptors
	Immediately before and 3 or 4h after MCA occlusion, i.p.	3 mg/kg, i.p.	Reduced the infarction volume induced by cerebral artery occlusion	Independent of CB <sub>1</sub> receptors
	Immediately before and 3h after MCA occlusion, i.p.	3 mg/kg, i.p.	Reduced the infarction volume induced by cerebral artery occlusion	5HT <sub>1A</sub> receptors dependent and TRPV <sub>1</sub> receptors independent
	Immediately before or 3h after MCA occlusion/ Repeated injections for 14 days and then before and 3h after MCA occlusion	0.1, 1 and 3 mg/kg, i.p.	Reduced the infarction volume induced by cerebral artery occlusion without development of tolerance	Independent of CB <sub>1</sub> and CB <sub>2</sub> receptors
6-hydroxydopamine toxicity <i>in vivo</i> and <i>in vitro</i>	2 weeks/daily	3 mg/kg, i.p.	Attenuated the reduction of tyrosine hydroxylase activity in the lesioned striatum and the reduction of this enzyme in the substantia nigra;	NS
amyloid $\beta$ -induced neuronal toxicity and microglial-conditioned media-based neurotoxicity <i>in vitro</i>	24-h incubation	10 $\mu$ M	Protected against A $\beta$ -evoked cell viability and from BV-2-conditioned media activated via LPS	NS
amyloid $\beta$ -induced toxicity and tert-butyl hydroperoxide-induced oxidative stress	15 min pre-incubation before A $\beta$ or sA $\beta$ addition/ 24-h incubation for oxidative stress analysis	0.01-10 $\mu$ M	Improved cell viability in response to tert-butyl hydroperoxide	NS
LPS-induced NO generation; microglial cell migration Morris water maze test in $\beta$ amyloid-injected mice,	24-h incubation 3 weeks: first week treated daily; second and third weeks treated 3 times/week, i.p.	10-1000 nM	Inhibited NO generation and ATP-induced intracellular calcium increase in cultured microglia; promoted microglial cell migration; prevented A $\beta$ -induced learning deficit and IL-6 mRNA expression	Some of the <i>in vitro</i> effects were mediated by A <sub>2A</sub> , CB <sub>1</sub> and CB <sub>2</sub> receptors
Experimental autoimmune encephalomyelitis (EAE)	Before anticipated relapse	5 and 10 mg/kg	Slowed the accumulation of disability from the inflammatory penumbra during relapsing EAE	Blockage of voltage-gated sodium channels

Paclitaxel (PAC)-induced mechanical sensitivity	Before each of the four PAC injections (On experimental days 1, 3, 5 and 7)	2.5, 5 and 10 mg/kg, i.p.	Prevented the PAC-induced mechanical sensitivity	5HT <sub>1A</sub> receptors dependent CB <sub>1</sub> and CB <sub>2</sub> receptors independent
N-metil-D-aspartate (NMDA)-induced retinal neurotoxicity	Immediately before intravitreal injection of NMDA (80 nmol/eye)	2 mg/kg, i.v.	Attenuated the NMDA-induced tyrosine nitration and reduced NMDA-induced apoptosis	Antioxidant: reduction of oxidative and nitrative stress
Diabetic retinopathy	Every 2 days for 2 or 4 weeks	10 mg/kg, i.p.	Reduced neural cell death, malondialdehyde (MDA) levels, 2'7'-dichlorofluorescein (DCF) fluorescence, tyrosine nitration, VEGF expression, ICAM-1 expression and TNF- $\alpha$ levels; blocked the increases of phosphorylation of p38	Antioxidant: reduction of oxidative and nitrative stress
amyloid $\beta$ -induced toxicity	24-h incubation	10 <sup>-7</sup> -10 <sup>-4</sup> M	Increased cell survival; decreased ROS production, MDA levels, caspase 3 levels and DNA fragmentation	Antioxidant: reduction of oxidative and nitrative stress
H <sub>2</sub> O <sub>2</sub> -induced oxidative stress	2-h incubation	1 $\mu$ M	Decreased cell death and DCF fluorescence	Antioxidant: reduction of oxidative stress
6-hydroxydopamine toxicity	2 weeks daily	3 mg/kg	Recovered 6-hydroxydopamine-induced dopamine depletion and upregulated mRNA levels of SOD in the substantia nigra	Antioxidant
Encephalopathy (bile duct ligation)	4 weeks	5mg/Kg	Improve of cognition and motor activity. Restores BDNF levels	5HT1A
Encephalopathy (thioacetamide)	Single dose	5mg/Kg	Cannabidiol restored liver function, normalizes 5-HT levels and improves brain pathology	5HT-dependent mechanism
$\beta$ amyloid-induced neurotoxicity	incubation	10 $\mu$ M	inhibited of phosphorylated form of p38 MAP kinase and transcription factor nuclear factor-kappaB activation.	Antioxidant
Pro-neurogenic	4 weeks	ND	Increased proliferation, survival and maturation of new neurons	CB1
Genetic model of Alzheimer's Disease	15 days	10mg/Kg	Decreased gliosis, neuronal death and facilitates neurogenesis	PPAR $\gamma$
amyloid $\beta$ -induced neurotoxicity	1 week	10mg/Kg	Decreased IL-1beta, GFAP and iNOS expression	NS
amyloid $\beta$ -induced neurotoxicity	24h	100nM	Decreased amyloid- $\beta$ production	PPAR $\gamma$
LPS-induced neurotoxicity	Single dose	3mg/Kg	blocked LPS-induced increase in TNF-alpha, COX-2 and blood brain barrier disruption.	NS
Pneumococcal meningitis	9 days	10mg/Kg	Prevented cognitive impairment, decreases TNF-alpha and IL6 in the brain.	NS
Amphetamine-induced oxidative stress	2 weeks	60mg/Kg	Decreased of carbonyl groups and prevents amphetamine-induced decreased BDNF expression.	Oxidative stress
Stress-reduced neurogenesis	2weeks 24h incubation	100n-30mg/Kg	Increased survival, differentiation and maturation	FAAH, CB1/CB2
NMDA- agonist induce psychosis	2 weeks	30-60mg/Kg	Decreased microglia activation	NS
LPS-induced microglia activation	2h	10 $\mu$ M	Blocked LPS-induced STAT1 activation	NF- $\kappa$ B and IFN $\beta$ -dependent pathways
encephalitogenic T cells	18h	5 $\mu$ M	Decreased Th17 activity, induction of CD4+CD25-CD69+LAG3+	NS

Cerebral malaria	7 days	30mg/Kg	Increased survival, reduce cognitive impairment, decrease IL-6 and TNF, increase BDNF	NS
------------------	--------	---------	---	----

NS: not studied.